

1

(a) (5 points) Halle el siguiente determinante:

$$\begin{vmatrix} 1 & 2 & 1 & 2 \\ 1 & 1 & 2 & 2 \\ 2 & 2 & 1 & 1 \\ 2 & 1 & 2 & 1 \end{vmatrix}$$

(b) (5 points) Sean  $A, B$  matrices cuadradas del mismo orden e invertibles. Halle la matriz  $X$  que soluciona el sistema matricial siguiente:

$$(A + B)(X + B) = B(X + A + B).$$

**Solución:**

(a) El determinante es equivalente a

$$\begin{vmatrix} 1 & 2 & 1 & 2 \\ 0 & -1 & 1 & 0 \\ 0 & -2 & -1 & -3 \\ 0 & -3 & 0 & -3 \end{vmatrix} = (-1) \begin{vmatrix} -1 & -3 \\ 0 & -3 \end{vmatrix} - \begin{vmatrix} -2 & -3 \\ -3 & -3 \end{vmatrix} = -3 - (6 - 9) = -3 + 3 = 0.$$

(b)  $X = A^{-1}(BA - AB)$ .



CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE  
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

---

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS  
CALL OR WHATSAPP: 689 45 44 70

[2]

Sea  $a \in \mathbb{R}$ . Se considera el sistema lineal

$$\begin{cases} x - 2y + z = 0 \\ -x + y + 2z = 2 \\ x - 3y + z = a \end{cases}$$

- (a) (5 points) Discuta el sistema.  
(b) (5 points) Resuelva el sistema. ¿Es  $(1, 2, 1)$  solución?

**Solución:**

- (a) La matriz  $A$  del sistema es cuadrada con determinante nulo. La matriz aumentada  $A^*$  es equivalente a

$$\left( \begin{array}{ccc|c} 1 & -2 & 1 & 0 \\ 0 & -1 & 1 & 0 \\ 0 & 0 & 0 & a-2 \end{array} \right).$$

Luego  $A^*$  tiene el mismo rango que  $A$  si y sólo si  $a = 2$ . Por tanto, el sistema no admite solución si  $a \neq 2$ . Cuando  $a = 2$ , el sistema es compatible indeterminado, dado que el rango de  $A$  es dos, menor que el número de variables. (Teorema de Rouché).

- (b) En el caso  $a = 2$ , utilizando el sistema escalonado equivalente hallado en el apartado anterior, es fácil ver que las soluciones son de la forma  $(5z - 4, 3z - 2, z)$ , donde  $z \in \mathbb{R}$ . Observamos que  $(1, 2, 1)$  no es solución, ya que  $z = 1$  corresponde con  $(1, 1, 1)$ , no con  $(1, 2, 1)$ .



CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE  
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

---

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS  
CALL OR WHATSAPP: 689 45 44 70

[3]

Se considera la función  $f(x) = \sqrt{\frac{4x-1}{x-4}}$ .

- (a) (5 points) Calcule el dominio y las asíntotas de  $f$ .  
(b) (5 points) Asumiendo que  $f$  admite inversa (no es necesario probarlo), halle la inversa de  $f$  y calcule su dominio.

**Solución:**

- (a) The domain is  $(4, \infty)$  and  $x = 4$  is a vertical asymptote since  $\lim_{x \rightarrow 4^+} f(x) = \infty$ . Since  $\lim_{x \rightarrow \infty} \frac{\sqrt{4x-1}}{\sqrt{x-4}} = 4$ , we have that  $y = 4$  is an horizontal asymptote. There is no oblique asymptote.
- (b) The inverse exists since it is asserted that the function is strictly monotonous. To find the inverse, we have to solve for  $x$  in the equation  $\sqrt{\frac{4x-1}{x-4}} = y$ . We get  $4x-1 = y^2(x-4)$ , that is,  $x = \frac{1-4y^2}{y^2-4}$ .



CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE  
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

---

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS  
CALL OR WHATSAPP: 689 45 44 70

[4]

- (a) (3 points) Demuestre que la ecuación  $x^2 - 2\sqrt{2x} + x - 1 = 0$  tiene al menos una solución positiva.
- (b) (7 points) Se considera la función  $f(x) = \frac{5x}{1+x^2}$ . Estudie los extremos locales y globales de  $f$  en el intervalo  $[0, 2]$ .

**Solución:**

- (a) The function  $g(x) = x^2 - 2\sqrt{2x} + x - 1$  is continuous in  $[0, \infty)$ . Note that  $g(0) = -1 < 0$  and  $g(2) = 1 > 0$ , thus by Bolzano's Theorem there is some  $0 < c < 2$  such that  $g(c) = 0$ , that is, there is  $c$  which is a positive solution of the proposed equation.
- (b) The function is continuous in  $[0, 2]$ , thus according to Weierstrass' Theorem,  $f$  has global maximum and global minimum in this interval. Moreover, it is derivable, thus we can locate possible extrema by equating the derivative of  $f$  to zero.

$$f'(x) = 5 \frac{1-x^2}{(1+x^2)^2} = 0 \text{ if and only if } x = \pm 1.$$

Now,  $-1 \notin [0, 2]$ , so it is disregarded. We have three candidates for being extremum of  $f$ : 1, 0 and 2. Evaluating we get  $f(1) = \frac{5}{2}$ ,  $f(0) = 0$  and  $f(2) = 2$ . Thus,  $x = 0$  is the global minimum and  $x = 2$  is the global maximum. Noting that  $f'$  is negative at the left of  $x = 1$  and positive at the right, we conclude that  $x = 1$  is local minimum.



CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE  
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

---

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS  
CALL OR WHATSAPP: 689 45 44 70

[5]

Sean  $a, b \in \mathbb{R}$ . Considere la función  $f(x) = \begin{cases} 1 + x^2, & \text{if } x \leq 0; \\ e^{1-2a+bx}, & \text{if } x > 0. \end{cases}$

- (a) (4 points) Estudie si  $f$  es continua en  $x = 0$ .  
(b) (6 points) Estudie si  $f$  es derivable en  $x = 0$ .

**Solución:**

- (a) Clearly the left sided limit of  $f$  at  $x = 0$  is 1, and the right sided limit is  $e^{1-2a}$ . So, in order to get continuity,  $a = \frac{1}{2}$ .
- (b) Since a differentiable function is also continuous,  $f$  is not differentiable at  $x = 0$  when  $a \neq \frac{1}{2}$ . Hence, suppose that  $a = \frac{1}{2}$ . After plugging this value into the expression of  $f$  we see that the left sided derivative is  $2x$  and the right sided derivative is  $be^{bx}$ . At  $x = 0$  these expressions take the value 0 and  $b$ , respectively. Thus,  $f$  is differentiable at  $x = 0$  if and only if  $a = \frac{1}{2}$  and  $b = 0$ .



CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE  
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

---

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS  
CALL OR WHATSAPP: 689 45 44 70

[6]

- (a) (5 points) Halle la integral de la función  $f(x) = x \ln x$ . (Pista: use integración por partes.)  
(b) (5 points) Halle el área de la región limitada por al gráfica de las funciones  $f(x) = x - 1$  y  $g(x) = -x^2 + x + 3$ .

**Solución:**

- (a) Take  $u = \ln x$  and  $dv = x dx$ . Then  $du = \frac{1}{x} dx$  and  $v = \frac{x^2}{2}$ . Apply the formula of integration by parts  $\int u dv = uv - \int v du$ , to get

$$\int x \ln x dx = \frac{x^2}{2} \ln x - \int \frac{x}{2} dx = \frac{x^2}{2} \ln x - \frac{x^2}{4} + C.$$

(b)  $A = \int_{-2}^2 (g(x) - f(x)) dx = \int_{-2}^2 4 - x^2 dx = 4x - \frac{x^3}{3} \Big|_{-2}^2 = \frac{32}{2} u^2.$



CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE  
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

---

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS  
CALL OR WHATSAPP: 689 45 44 70