

Functions several variables $f: \mathbb{R}^N \rightarrow \mathbb{R}^M$

$\text{Dom} f \equiv$ points where the function is well-defined

$\text{Im} f \equiv$ points $y \in \mathbb{R}^M$ such that there exists $x \in \mathbb{R}^N$ with $f(x) = y$

Example: Problem 8 (i)

$$f(x, y) = \frac{1}{xy}$$

$$f: \mathbb{R}^2 \rightarrow \mathbb{R}$$

\uparrow \uparrow
Dom f Im f .

$$\text{Dom} f = \{ (x, y) \in \mathbb{R}^2 ; xy \neq 0 \}$$

$$= \{ (x, y) \in \mathbb{R}^2 ; x \neq 0 \text{ or } y \neq 0 \} \subset \mathbb{R}^2$$

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Type of functions

Scalar functions

$f: \mathbb{R}^N \rightarrow \mathbb{R}$ the image is a number

Vector functions

$F: \mathbb{R}^N \rightarrow \mathbb{R}^M$ the image is a vector

In particular, vector fields

if $N = M$

$F: \mathbb{R}^N \rightarrow \mathbb{R}^N$

Examples:

Parametric equations of a line in \mathbb{R}^3

$$\left. \begin{aligned} x &= t \\ y &= 2 + 3t \\ z &= 1 + t \end{aligned} \right\}$$

$$F(x, y, z) = \begin{pmatrix} 0 \\ 2 \\ 1 \end{pmatrix} + t \begin{pmatrix} 1 \\ 3 \\ 1 \end{pmatrix} \in \mathbb{R}^3$$

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Secret.

$$F(x, y, z) = (xy, x^2y, \log x)$$

scalar functions \equiv magnitudes
vector fields \equiv velocity, etc.

Definition - Level curves

Let $A \subset \mathbb{R}^n$ be a subset and $f: A \rightarrow \mathbb{R}$

The level set of f at the value c as
the set of points $x \in A$ such that

$$f(x) = c$$

\curvearrowright TO $N=2$ ($\text{Dom} f \subset \mathbb{R}^2$) we find level curves


CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Cartagena99

Recall

- The level curves are points of the domain where the function is constant.
- It allows us to draw 3D figures in 2D



Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70


of new quality the function curves

Problem 9: $f(x,y) = xy$, $c = 1, -1, 3$

$$c = 1 \Rightarrow xy = 1 \quad ; \quad y = \frac{1}{x}$$

$$c = -1 \Rightarrow xy = -1 \quad ; \quad y = -\frac{1}{x}$$

$$c = 3 \Rightarrow xy = 3 \quad ; \quad y = \frac{3}{x}$$



Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Example: Describe the graph of
 $f(x,y) = x^2 + y^2$ $f \equiv$ function

$f: \mathbb{R}^2 \rightarrow \mathbb{R}$. $\text{Dom} f = \mathbb{R}^2$

$f \geq 0$


Vertical planes.

$x=0 \Rightarrow f(0,y) = y^2$

$x=x_0$




$f(x_0,y) = y^2 + x_0^2$



CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
 LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
 CALL OR WHATSAPP: 689 45 44 70

Cartagena99



Limits and continuity

The limit of a function at a point x_0 will be the value that the function should be depending on the values around it.

$$\lim_{x \rightarrow x_0} f(x) = L$$

Definition $f: \mathbb{R}^N \rightarrow \mathbb{R}^M$, $x_0 \in \mathbb{R}^N$, $L \in \mathbb{R}^M$

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Cartagena99

How to compute the limits?

① Applying the definition.

For any $\varepsilon > 0$, $\exists \delta > 0$, $\delta = \delta(\varepsilon)$
such that (for example in 2D) $f: \mathbb{R}^2 \rightarrow \mathbb{R}$.

$$0 < \|(x, y) - (x_0, y_0)\| = \sqrt{(x-x_0)^2 + (y-y_0)^2} < \delta$$

$$0 < (x-x_0)^2 + (y-y_0)^2 < \delta^2$$



$$|f(x, y) - L| < \varepsilon \quad \leftarrow \text{we start from here}$$

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Example: $\lim_{(x,y) \rightarrow (0,0)} \sqrt{x^2+y^2} \sin \frac{1}{x^2+y^2} = 0$

Find $\delta > 0$, we start from

$$|f(x,y) - 0| = |f(x,y)| = \left| \sqrt{x^2+y^2} \sin \frac{1}{x^2+y^2} \right|$$

$$\leq \sqrt{x^2+y^2} < \epsilon = \delta$$

$$\delta = \delta(\epsilon) = \epsilon$$

and choosing those δ and ϵ , when (x,y) are suff. close to $(0,0)$ the function $f(x,y)$ will be suff. close to 0.

□

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Cartagena99

② Iterative limits

$$\lim_{x \rightarrow a} \left(\lim_{y \rightarrow b} f(x, y) \right) = \lim_{y \rightarrow b} \left(\lim_{x \rightarrow a} f(x, y) \right) = L$$

This method is valid to prove non-existence.
but not to prove the value of the limit.

③ Directional limits . Approximation through a family of functions

For example: $\left. \begin{array}{l} y = \lambda x \\ y = \lambda x^2 \\ \vdots \end{array} \right\}$


Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Remark

In one variable we used to compute one-sided limits to prove existence of a limit.



In several variables we have infinitely many directions so the approach following a family of functions does not justify the existence of the limit.

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70


ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Example: $\lim_{(x,y) \rightarrow (0,0)} \frac{x}{\sqrt{x^2+y^2}}$

We choose the family of curves $y = \lambda x$,
 $\lambda \equiv$ arbitrary parameter.

Then,

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x}{\sqrt{x^2+y^2}} = \lim_{x \rightarrow 0} \frac{x}{\sqrt{x^2+\lambda^2 x^2}} = \lim_{x \rightarrow 0} \frac{\pm 1}{\sqrt{1+\lambda^2}}$$



$$= \frac{x}{|x| \sqrt{1+\lambda^2}} = \frac{\pm 1}{\sqrt{1+\lambda^2}}$$

The limit depends on λ , on the direction
 in which the limit does not exist.

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
 LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
 CALL OR WHATSAPP: 689 45 44 70

Problem 16

$$f(x,y) = \begin{cases} \frac{x^3 y}{x^6 + y^2} & \text{if } (x,y) \neq (0,0) \\ 0 & \text{if } (x,y) = (0,0) \end{cases}$$

i) $\lim_{(x,y) \rightarrow (0,0)} f(x,y)$ along $y = x$

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^3 y}{x^6 + y^2} = \lim_{x \rightarrow 0} \frac{x^4}{x^6 + x^2}$$

$= \lim_{x \rightarrow 0} \frac{x^2}{x^4 + 1} = 0 \Rightarrow$ The limit following $y = x$ is 0 but it does not mean the limit exists.

... to limit along $y = x^3$




CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

exist.

④ limits using polar coordinates (in \mathbb{R}^2)

$$\begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases}$$



Proposition

Let A be a subset in \mathbb{R}^2 such that $(0,0) \in A$

and $f: A \subset \mathbb{R}^2 \rightarrow \mathbb{R}$

If $\lim_{r \rightarrow 0} f(r \cos \theta, r \sin \theta)$

depends on θ , then $\lim_{(x,y) \rightarrow (0,0)} f(x,y)$ does not exist.

For a general point $(x_0, y_0) \in A$ this is valid

" " " " +

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

$(x,y) \rightarrow (x_0, y_0)$

Cartagena99

Proposition

$A \subset \mathbb{R}^N$, $x_0 \in A$, $f, g : A \subset \mathbb{R}^N \rightarrow \mathbb{R}$
two scalar functions.

If $\lim_{x \rightarrow x_0} f(x) = 0$ and g is bounded
 $|g(x)| < C, x \in B(x_0, \delta)$

Then,
 $\lim_{x \rightarrow x_0} f(x)g(x) = 0$

Example: $\lim_{(x,y) \rightarrow (0,0)} \frac{xy^2}{x^2+y^2} =$


Assume $\left. \begin{array}{l} x = r \cos \theta \\ y = r \sin \theta \end{array} \right\} \begin{array}{l} \rightarrow 0 \\ \rightarrow 0 \end{array}$

Cartagena99


CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

$$= \lim_{\theta \rightarrow 0} 2 \underbrace{\cos \theta \sin^2 \theta}_{\text{bounded}} = 0$$



Therefore we can conclude that the limit exists and it is actually 0.



Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Continuity

Definition

A subset of \mathbb{R}^N , $x_0 \in \Delta$, $f: \Delta \subset \mathbb{R}^N \rightarrow \mathbb{R}^M$
we say that f is continuous if

$$\lim_{x \rightarrow x_0} f(x) = f(x_0)$$

Assuming:

- f is defined at x_0
- The limit exists
- $\lim_{x \rightarrow x_0} f(x) = f(x_0)$

In case $f(x) = (f_1(x), \dots, f_m(x))$
it will be cont. if every

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Cartagena99

Example:

$$f(x,y) = x^2 \sin(x^2 + y)$$

cont. since it is a composition, addition and multiplication of cont. functions.

$$F(x,y) = \left(x^2 y \sin(x^2 + y), \frac{x}{x^2 + y^2 + 1} \right)$$

Both components are cont. since $x^2 + y^2 + 1 \neq 0$ for any $(x,y) \in \text{Dom } F$.

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70