

Derivations: Partial derivations


$$\text{If } f(x,y), \quad \frac{\partial f(x,y)}{\partial x}, \quad \frac{\partial f(x,y)}{\partial y}$$

ratio of change on the direction of the axis.

Directional derivations:

$$D_{\vec{v}} f(x,y), \quad \vec{v} = \text{vector.}$$

ratio of change on the direction \vec{v} .



CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Cartagena99

Differentiability

We have $f: \mathbb{R}^2 \rightarrow \mathbb{R}$ a scalar function
and we would like to see if


f is differentiable at a point P .

17:25 Sun 22 Mar

83%

general function.pdf

📄 🗨️ ⋮



Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

At a point $(x_0, y_0, f(x_0, y_0)) \in \mathbb{R}^3$ will have
a plane going through that point with equation

any
plane. $\left\{ \begin{array}{l} z = f(x_0, y_0) + A(x - x_0) + B(y - y_0) \end{array} \right.$

$A, B \in \mathbb{R}$ coefficients for the
plane.

But, if we want to have the tangent plane
to the surface $z = f(x, y)$ at $P = (x_0, y_0, f(x_0, y_0))$

We need that

$$f\left(x_0, y_0 + \begin{array}{c} (x-x_0) \\ \hline \end{array}, \begin{array}{c} (y-y_0) \\ \hline \end{array}\right) = f(x_0, y_0) + A h + B k + \underline{\underline{2(h, k)}}$$

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

$h-x_0$

That would be the tangent plane if

$$\lim_{(h,k) \rightarrow (0,0)} \frac{z(h,k)}{\|(h,k)\|} = 0 = \lim_{(x,y) \rightarrow (x_0,y_0)} \frac{z((x,y)-(x_0,y_0))}{\|(x,y)-(x_0,y_0)\|}$$

Definition - Differentiability

Let $A \subset \mathbb{R}^2$ be a set in \mathbb{R}^2 such that $(x_0, y_0) \in A$

$f: A \subset \mathbb{R}^2 \rightarrow \mathbb{R}$ a scalar function

Then, f is differentiable at (x_0, y_0) if

a) $\frac{\partial f(x_0, y_0)}{\partial x}, \frac{\partial f(x_0, y_0)}{\partial y}$ exist

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Cartagena99

We might write that limit as

$$\lim_{(x,y) \rightarrow (x_0,y_0)} \frac{f(x,y) - f(x_0,y_0) - \nabla f(x_0,y_0) \cdot \begin{pmatrix} x-x_0 \\ y-y_0 \end{pmatrix}}{\|(x,y) - (x_0,y_0)\|} = 0$$

If that is the case we write the tangent plane to the graph of f at (x_0, y_0) as

$$z = f(x_0, y_0) + \underbrace{\frac{\partial f(x_0, y_0)}{\partial x}}_A (x - x_0) + \underbrace{\frac{\partial f(x_0, y_0)}{\partial y}}_B (y - y_0)$$

These two particular coefficients

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Cartagena99


Definition

$\Delta \subset \mathbb{R}^N$, $x_0 \in \Delta$, $f: \Delta \rightarrow \mathbb{R}^M$

f is differentiable at $x_0 \in \mathbb{R}^N$ if

a) all partial derivatives exist at x_0
Jacobian matrix → *recta*

b) $\lim_{x \rightarrow x_0} \frac{\|f(x) - f(x_0) - Jf(x_0)(x - x_0)\|}{\|x - x_0\|} = 0$



Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Problem 5 of set 1.2

$$f(x,y) = \begin{cases} \frac{2xy}{x^2+y^2} & \text{if } (x,y) \neq (0,0) \\ 0 & \text{if } (x,y) = (0,0) \end{cases}$$

c) f differentiable at $(0,0)$?

We might study the continuity first at $(0,0)$

$$\lim_{(x,y) \rightarrow (0,0)} f(x,y) = \lim_{(x,y) \rightarrow (0,0)} \frac{2xy}{x^2+y^2} = \lim_{x \rightarrow 0} \frac{2 \cdot x^2}{x^2(1+1^2)}$$

$$y = 2x$$

$$= \lim_{x \rightarrow 0} \frac{2x}{1+1^2} = \frac{2x}{1+1^2}$$

The limit depends on the direction so it

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

$$\lim_{(x,y) \rightarrow (0,0)} \frac{2xy}{x^2+y^2} = \lim_{r \rightarrow 0} \frac{2r^2 \cos\theta \sin\theta}{r^2} = 2 \cos\theta \sin\theta$$

$$\begin{cases} x = 2 \cos\theta \\ y = 2 \sin\theta \end{cases}$$

The limit depends on the direction θ so it does not exist.

Existence of all directional derivatives $\not\Rightarrow$ continuity

\neq differentiability.

ii) Find $\frac{\partial f}{\partial x}$, $\frac{\partial f}{\partial y}$ at $(0,0)$

Show that $\frac{\partial f}{\partial x}$ is not continuous.

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

$$\lim_{h \rightarrow 0} \frac{f(h, 0) - f(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{0 - 0}{h} = 0$$

$$f(x, y) = \frac{2xy}{x^2 + y^2} \Rightarrow f(h, 0) = \frac{2 \cdot 0 \cdot h}{h^2 + 0} = 0$$

$$f(0, 0) = 0$$

$$f(0, k) = \frac{2 \cdot 0 \cdot k}{0 + k^2} = 0$$

Partial derivative with respect to x

If $(x, y) \neq (0, 0)$ (rules of derivatives)

$$\frac{\partial f(x, y)}{\partial x} = \frac{2y(x^2 + y^2) - \frac{4x^2y}{2x}}{(x^2 + y^2)^2}$$

2x2y2y3

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Cartagena99

$$\frac{\partial f(x,y)}{\partial x} = \begin{cases} \frac{-2x^2y + 2y^3}{(x^2+y^2)^2} & \text{if } (x,y) \neq (0,0) \\ 0 & \text{if } (x,y) = (0,0) \end{cases}$$

Continuous at (0,0) if

$$\lim_{(x,y) \rightarrow (0,0)} \frac{-2x^2y + 2y^3}{(x^2+y^2)^2} = 0 = \frac{\partial f(0,0)}{\partial x}$$

Using polar coordinates. $\begin{cases} x = r \cos \theta \\ y = r \sin \theta \end{cases}$

$$\lim_{r \rightarrow 0} \frac{-2x^2y + 2y^3}{(x^2+y^2)^2} = \lim_{r \rightarrow 0} \frac{-2r^3 \cos^2 \theta \sin \theta + 2r^3 \sin^3 \theta}{r^4}$$

$= \infty$ bounded.

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Problem 10 i)

$$\left. \begin{aligned} f(x, y) &= x - y + 2 \\ (x_0, y_0) &= (1, 3) \end{aligned} \right\} \text{Find the tangent plane.}$$

f linear so that tangent plane $\equiv f$.

$$z = f(x_0, y_0) + \frac{\partial f(x_0, y_0)}{\partial x}(x - x_0) + \frac{\partial f(x_0, y_0)}{\partial y}(y - y_0)$$


$$f(x_0, y_0) = 1 - 3 + 2 = 0$$

$$\frac{\partial f(x_0, y_0)}{\partial x} = 1 \quad \frac{\partial f(x_0, y_0)}{\partial y} = -1$$


Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70



But f is away from T if we are outside of the ball



CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP:689 45 44 70

Cartagena99

Proposition

$\Lambda \subset \mathbb{R}^N$, $x_0 \in \Lambda$ and $f: \Lambda \rightarrow \mathbb{R}$ differentiable at x_0 and $v \in \mathbb{R}^N$ for a vector.

Then,

$$D_v f(x_0) = \sum_{i=1}^N \frac{\partial f(x_0)}{\partial x_i} \cdot v_i = \langle \nabla f(x_0), v \rangle$$

v is normalised vector

$$\|v\| = 1$$

$$\bullet \langle \nabla f(x_0), (\alpha v) \rangle = \alpha \langle \nabla f(x_0), v \rangle$$

\neq if $\alpha \neq 1$

$$\langle \nabla f(x_0), v \rangle$$

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Example: Set 1.3 problem ① i)

$f(x,y) = x^2 + y^2$ at $(1,1)$ along the direction $(1,-1)$

$$D_{\vec{v}} f(1,1) = \lim_{(x,y) \rightarrow (1,1)} \frac{f((1,1) + t(1,-1)) - f(1,1)}{t \|(1,-1)\|}$$

f is differentiable everywhere (it is a polynomial)

$$D_{\vec{v}} f(1,1) = \left\langle \nabla f(1,1), \frac{(1,-1)}{\|(1,-1)\|} \right\rangle$$

$$\nabla f(x,y) = (2x, 2y) \Rightarrow \nabla f(1,1) = (2, 2)$$

$$\|(1,-1)\| = \sqrt{1+1} = \sqrt{2}$$

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Remark

$$-\frac{\|\nabla f(x_0)\|}{\|\nabla f(x_0)\|} \nabla_{\sigma} f(x_0) = \langle \nabla f(x_0), \sigma \rangle = \|\nabla f(x_0)\| \underbrace{\|\sigma\|}_{=1} \cos(\nabla f, \sigma)$$

$$= \|\nabla f(x_0)\| \cos(\nabla f, \sigma) \leq \|\nabla f(x_0)\|$$

We obtain that the directional derivative of f at x_0 in the direction of σ is maximal in the direction of ∇f

$$\cos(\nabla f, \sigma) = 1 \Rightarrow \text{angle}(\nabla f, \sigma) = 0$$

σ

$\nabla f / \|\nabla f\|$

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

- Problem 5, set 1.3

Temperature of a metal plate

$$T(x,y) = e^x \cos y + e^y \cos x$$

- a) Direction of maximal increasing for T at $(0,0)$

$$\nabla T(x,y) = \left(\underbrace{e^x \cos y - e^y \sin x}_{\frac{\partial T}{\partial x}}, \underbrace{-e^x \sin y + e^y \cos x}_{\frac{\partial T}{\partial y}} \right)$$

$$\nabla T(0,0) = (1, 1)$$

- b) T decreasing the fastest

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Proposition

$\Delta \subset \mathbb{R}^n$, $x_0 \in \Delta$, $f: \Delta \rightarrow \mathbb{R}$ differentiable at x_0
with $\nabla f(x_0) \neq 0$ then
 $\nabla f(x_0) \perp$ to the level curve of f at $f(x_0)$


For example in \mathbb{R}^3

$$\nabla f(x, y, z) \neq (0, 0, 0)$$

$(x, y, z) \in \Omega$, Ω domain in \mathbb{R}^3

Assume a level curve

$$f(x, y, z) = c \in \mathbb{R}$$




Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Since f is diff. the tangent plane will be perpendicular to the level curve.



so that ∇f is the

\vec{z} tangent

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Cartagena99

In other words, take two points on the tangent plane.

$$P^* = (P_1, P_2, P_3) \text{ and } P = (x, y, z)$$

with $n = (n_x, n_y, n_z)$ as the normal vector to the tangent plane.

$$(P - P^*) \cdot n = 0 \Rightarrow (x - P_1)n_x + (y - P_2)n_y + (z - P_3)n_z = 0$$

There is only one normal vector !!

$$\nabla f(x, y, z) \cdot (P - P^*) = 0$$

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70

Problem 11. Set 1.2 i)

Sphere.
Surface: $x^2 + y^2 + z^2 = 3$ at $(x_0, y_0, z_0) = (1, 1, 1)$

$$\nabla f(x, y, z) = (2x, 2y, 2z)$$

$$\nabla f(1, 1, 1) = (2, 2, 2)$$

Tangent plane:

$$\nabla f(1, 1, 1) \cdot \begin{pmatrix} x-1 \\ y-1 \\ z-1 \end{pmatrix} = 0$$

$$2x - 2 + 2y - 2 + 2z - 2 = 0$$

Cartagena99

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS
CALL OR WHATSAPP: 689 45 44 70