

## Problem 1

(a) Generate a vector  $\mathbf{x}$  containing 1000 independent replicates of a  $N(10, 4^2)$  random variable. Draw a histogram of  $\mathbf{x}$  using the `prob=TRUE` option. I suggest that you set `breaks=20` in the `hist` command — see `help(hist)` for an explanation of what this does.

Superimpose the probability density function of the  $N(10, 4^2)$  distribution on your histogram.

(b) Generate  $\mathbf{y1}$  containing 400 realisations of a  $N(3, 3^2)$  random variable,  $\mathbf{y2}$  containing 400 realisations of a  $N(5, 4^2)$  RV, and  $\mathbf{y3}$  containing 400 realisations of a  $N(7, 5^2)$  RV, all variables being independent. Calculate  $\mathbf{w}=\mathbf{y1}+\mathbf{y2}+\mathbf{y3}$  and draw a histogram of  $\mathbf{w}$ , using the `breaks` option to get a suitable number of bars in the histogram.

(c) Guess the distribution of  $W = Y_1 + Y_2 + Y_3$  when  $Y_1 \sim N(3, 9)$ ,  $Y_2 \sim N(5, 16)$  and  $Y_3 \sim N(7, 25)$  are independent RVs.

Re-draw the histogram of  $\mathbf{w}$  using the `prob=TRUE` option and superimpose the probability density function for your chosen distribution. Remember, in the `curve` command you have to give a function of  $x$  — not  $w$  — to be plotted (see “Plotting commands” on page 1).

(d) Carry out further simulations to give a more definitive check that your chosen PDF really is correct.

**In your submission:** In a Word document, include the content of the R console window showing R commands and the resulting output. You should also include the histograms of  $\mathbf{x}$  and  $\mathbf{w}$  from the graphics window and an explanation for your choice of distribution in part (c).

See “Graphics”, “Writing a plot directly to a file” and “Combining material in your coursework solutions” in the **Brief Introduction to R**.

## Problem 2

Hermione and Ron have won prizes with random values. Hermione’s prize will be  $H = 100 \exp(X)$  Sickles and Ron’s prize will be  $R = 100 \exp(Y)$  Sickles, where  $X$  and  $Y$  are independent  $N(0, 0.3^2)$  random variables.

(a) Write R commands to simulate one pair of values of  $(H, R)$  and compute  $W = H/R$ . Create a loop to run the above commands 200 times. Store the 200 values of  $H$  in a vector  $H_{sample}$ , the 200 values of  $R$  in

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE  
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

---

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS  
CALL OR WHATSAPP:689 45 44 70

ω √ J.036 π

Cartagena99

Investigate Professor Slughorn's claim by superimposing the above density on a histogram of *Wsample*. In order to obtain a useful plot, you may find it necessary to restrict attention to values of *W* below an upper limit, such as 5. You can do this with the command

```
hist(Wsample[Wsample<5],prob=TRUE).
```

Do your results support the Professor's theory?

(c)

Professor Slughorn explains that the CDF corresponding to the PDF (1) is

$$F_W(w) = \Phi\left(\frac{\log(w)}{\sqrt{0.18}}\right)$$

where  $\Phi$  is the standard normal CDF.

Evaluate the function  $F_W(w)$  at  $w \in \{0.2, 0.4, 0.6, 0.8, 1, 1.2, 1.4, 1.6, 1.8, 2\}$ . Find the proportions of values in *Wsample* less than each of these values of  $w$  and plot the proportions against  $F_W(0.2), F_W(0.4), \dots, F_W(1.8), F_W(2)$ . What does this plot show?

(d) Repeat the comparison conducted in part (c) for larger data sets and state whether you believe Professor Slughorn's claim is correct.

**In your submission:** Give the content of the R console window; include the relevant plots; give clear written answers in parts (b), (c) and (d).

The logo for Cartagena99 features the text 'Cartagena99' in a stylized, blue, serif font. The '99' is significantly larger and more prominent than the rest of the text. The logo is set against a light blue background with a white starburst shape behind the text.

CLASES PARTICULARES, TUTORÍAS TÉCNICAS ONLINE  
LLAMA O ENVÍA WHATSAPP: 689 45 44 70

---

ONLINE PRIVATE LESSONS FOR SCIENCE STUDENTS  
CALL OR WHATSAPP:689 45 44 70